

Pharmacological management of asthma in children aged 5–11 years

The recently updated Asthma and Respiratory Foundation NZ guidelines (2020) did not contain any major changes to the diagnosis or management of asthma in children aged 5–11 years. A SABA reliever used as needed (without an inhaled corticosteroid [ICS]) continues to be recommended in children who are on Step 1 (Figure 1). This is in contrast to the updated guidelines for adolescents and adults, which no longer recommend use of SABA alone (to be covered in a future article).²

Stepwise treatment of asthma in children aged 5–11 years¹

Pharmacological management of asthma in children follows a stepwise progression where stepping treatment up or down is guided by symptom control and risk of exacerbations.¹ Before stepping up, check inhaler technique (including use of a spacer), adherence, understanding of the management plan and any barriers to its implementation.¹ Once symptoms have been well-controlled for at least eight weeks, consider stepping down and reassess control after 12 weeks.¹

Step 1 is for children with intermittent symptoms and involves as needed use of a SABA, without maintenance treatment.

Step 2 is for children who have symptoms >2 times/week, use a reliever >2 times/week, had regular night waking with symptoms in the past month or had an exacerbation requiring oral corticosteroids in the past year. This step involves the addition of a daily low dose ICS (see Table 1 for dose definitions).

Montelukast, a leukotriene receptor antagonist, may be considered as an alternative to ICS, but ICS are generally more effective.

Step 3 is for children with uncontrolled symptoms despite optimal treatment at Step 2. This step involves the addition of a LABA in combination with an ICS. LABAs (with ICS) should only be initiated when the child is clinically stable. If the LABA is ineffective or symptoms worsen after starting it, treatment should be stopped.

Step 4 is for children with uncontrolled symptoms despite optimal treatment at Step 3. This step involves an increase from low dose ICS/LABA to standard dose ICS/LABA (Table 1). Montelukast may be added if control remains poor. If control at Step 4 is poor, refer for paediatric assessment.

Step 5 is for children with uncontrolled symptoms despite optimal treatment at Step 4. This step involves standard dose ICS/LABA and montelukast, if the child is not already taking this. High dose ICS/LABA may be considered (Table 1). The child should be referred for paediatric assessment.

N.B. An anti-inflammatory reliever (AIR) regimen that includes maintenance budesonide/formoterol (also known as Single combination ICS/LABA inhaler Maintenance And Reliever Therapy [SMART]) may be considered for children who are poorly controlled on Steps 3–5 on advice from a paediatric respiratory specialist.¹ AIR therapy/SMART is unapproved in children.






Figure 1. Stepwise treatment for managing asthma in children aged 5–11 years. Adapted from the Asthma and Respiratory Foundation NZ guidelines (2020).¹ See Table 1 for dose definitions. N.B. “High” dose is double the standard dose.

Oral corticosteroids are indicated for moderate and severe acute asthma

Initial treatment with SABA, delivered via a metered dose inhaler and spacer, and oral corticosteroids is likely to be sufficient for most children presenting with acute moderate or severe asthma.¹ While the evidence of benefit of oral corticosteroid treatment in children aged under five years is limited, in older children with acute asthma, oral corticosteroids reduce the need for hospital admission and the risk of relapse.³ If required, give oral prednisolone (liquid) or prednisone (tablet) at 1–2 mg/kg per day, up to maximum of 40 mg daily, for 3–5 days.¹ Tapering before stopping treatment is not necessary for short courses. Some asthma action plans for school-age children may include directions for when to initiate a course of oral corticosteroids prescribed for use at home. However, children who require frequent or continuous use of oral corticosteroids (more than 14 days in 12 months) should be referred for paediatric assessment.¹

 The Asthma and Respiratory Foundation NZ Child Asthma Guidelines 2020 are available from: <https://www.nzrespiratoryguidelines.co.nz/childguidelines-654716.html>

 An upcoming article will discuss the changes to the pharmacological management of asthma in adolescents and adults in the updated Asthma and Respiratory Foundation NZ guidelines.

References

1. Asthma and Respiratory Foundation NZ. New Zealand child asthma guidelines: a quick reference guide. 2020. Available from: https://www.nzrespiratoryguidelines.co.nz/uploads/8/3/0/1/83014052/arf_nz_child_asthma_guidelines_update_30.6.20.pdf (Accessed July, 2020).
2. Beasley R, Beckert L, Fingleton J, et al. NZ adolescent & adult asthma guidelines. NZMJ 2020;133. Available from: www.nzma.org.nz/journal-articles/asthma-and-respiratory-foundation-nz-adolescent-and-adult-asthma-guidelines-2020-a-quick-reference-guide (Accessed Jun, 2020)
3. National Asthma Council. Australian asthma handbook: Starting systemic corticosteroid treatment. 2019. Available from: <https://www.astmahandbook.org.au/acute-asthma/clinical/corticosteroids#evref-9> (Accessed Jul, 2020).

Table 1. Low and standard ICS doses for children recommended in the Asthma and Respiratory Foundation NZ guidelines.

ICS (funded brand name)	Low dose	Standard dose
Beclomethasone dipropionate (Beclazone)	200 micrograms/day Prescribe: ■ Beclazone 100, one actuation, twice daily OR ■ Beclazone 50, two actuations, twice daily	400–500 micrograms/day Prescribe: ■ Beclazone 100, two actuations, twice daily OR ■ Beclazone 250, one actuation, twice daily
Beclomethasone dipropionate ultrafine (Qvar)	100 micrograms/day Prescribe: ■ Qvar 50, one actuation, twice daily	200 micrograms/day Prescribe: ■ Qvar 100, one actuation, twice daily
Budesonide (Pulmicort)	200 micrograms/day Prescribe: ■ Pulmicort 100, one actuation twice daily	400 micrograms/day Prescribe: ■ Pulmicort 200, one actuation, twice daily
Fluticasone propionate (Flixotide*, Floair†)	100 micrograms/day Prescribe: ■ Flixotide 50*, one actuation, twice daily OR ■ Floair 50†, one actuation, twice daily	200–250 micrograms/day Prescribe: ■ Flixotide 50*, two actuations, twice daily OR ■ Flixotide 125*, one actuation, twice daily OR ■ Floair 50†, two actuations, twice daily OR Floair 125†, one actuation, twice daily

* Prescribe metered dose inhaler (MDI) unless child is able to use the dry powder inhaler (accuhaler)

† From 1 September, 2020, the Floair brand of fluticasone MDI will no longer be funded. For further information, see: www.pharmac.govt.nz/news/notification-2019-12-12-fluticasone